

## Ku-70 (Acetyl Lys539) rabbit pAb

Cat No.: ES1107

For research use only

## Overview

Product Name Ku-70 (Acetyl Lys539) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300

ELISA: 1/20000. Not yet tested in other applications.

**Immunogen** Synthesized acetyl-peptide derived from the human

Ku-70 around the acetylation site of K539.

**Specificity** Acetyl-Ku-70 (K539) Polyclonal Antibody detects

endogenous levels of Ku-70 protein only when

acetylated at K539.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at  $-20^{\circ}$ C. Avoid repeated freeze-thaw cycles.

Protein Name X-ray repair cross-complementing protein 6

Gene Name XRCC6

**Cellular localization** Nucleus . Chromosome .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 70kD
Human Gene ID 2547
Human Swiss-Prot Number P12956

Alternative Names XRCC6; G22P1; X-ray repair cross-complementing

protein 6; 5'-deoxyribose-5-phosphate lyase Ku70; 5'-dRP lyase Ku70; 70 kDa subunit of Ku antigen;

ATP-dependent DNA helicase 2 subunit 1;

ATP-dependent DNA helicase II 70 kDa subunit;CTC

box-binding factor 7

**Background** The p70/p80 autoantigen is a nuclear complex

consisting of two subunits with molecular masses of



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus. [provided by RefSeq, Jul 2008],



+86-27-59760950