

NFKB-p65 (Acetyl Lys314/Lys315) rabbit pAb

Cat No.: ES1125

For research use only

Overview

Immunogen

Product Name NFκB-p65 (Acetyl Lys314/Lys315) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300

ELISA: 1/20000. Not yet tested in other applications. Synthesized acetyl-peptide derived from the human

NFκB-p65 around the acetylation site of K314/K315.

Specificity Acetyl-NFkB-p65 (K314/K315) Polyclonal

AntibodySynthesized peptide derived from the human NFkB-p65 around the acetylation site of

K314/K315.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Transcription factor p65

Gene Name RELA

Cellular localization Nucleus . Cytoplasm . Nuclear, but also found in the

cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B) (PubMed:1493333). Colocalized with DDX1 in the nucleus upon TNF-alpha induction (PubMed:19058135). Colocalizes with GFI1 in the nucleus after LPS stimulation (PubMed:20547752).

Translocation to the nucleus is impaired in

L.monocytogenes infection (PubMed:20855622). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 60kD
Human Gene ID 5970
Human Swiss-Prot Number Q04206




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com




Alternative Names

Background

RELA; NFKB3; Transcription factor p65; Nuclear factor NF-kappa-B p65 subunit; Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011],



Western Blot analysis of HeLa, HepG2 cells using Acetyl-NFκB-p65 (K314/K315) Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Western Blot analysis of Hela cells using Acetyl-NFκB-p65 (K314/K315) Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100

