

## Bad (phospho Ser136) rabbit pAb

## Cat No.:ES1270

For research use only

## Overview

Product Name	Bad (phospho Ser136) rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human; Mouse; Rat
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
_	peptide derived from human BAD around the
	phosphorylation site of Ser136. AA range:102-151
Specificity	Phospho-Bad (S136) Polyclonal Antibody detects
	endogenous levels of Bad protein only when
	phosphorylated at S136.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Bcl2 antagonist of cell death
Gene Name	BAD
Cellular localization	Mitochondrion outer membrane. Cytoplasm .
	Colocalizes with HIF3A in the cytoplasm (By
	similarity). Upon phosphorylation, locates to the
	cytoplasm
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	
Human Gene ID	572
Human Swiss-Prot Number	Q92934
Alternative Names	BAD; BBC6; BCL2L8; Bcl2 antagonist of cell death;
	BAD; Bcl-2-binding component 6; Bcl-2-like protein
	8; Bcl2-L-8; Bcl-XL/Bcl-2-associated death promoter



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C




Background

The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq, Jul 2008],

Western Blot analysis of various cells using Phospho-Bad (S136) Polyclonal Antibody

-		BAD 22KD	BAD 22KD
-		BAD (p-Ser136) 22KD	ł.
i.	+	- phospho-peptide	
-		+ non-phospho-peptide	
+	+	+ NIH/3T3 Insulin 0.01U/ml 15'	

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using BAD (Phospho-Ser136) Antibody. The picture on the right is blocked with the phospho peptide.





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C