

Recombinant Human FGFb (146AA)

| Catalog # | EPT078 |
|-----------------|---|
| Expression Host | E.coli |
| DESCRIPTION | Recombinant Human Fibroblast Growth Factor |
| | 2/Fibroblast Growth Factor Basic is produced by our |
| | E.coli expression system and the target gene encoding |
| | Pro143-Ser288 is expressed. |
| Accession | P09038 |
| Synonyms | Fibroblast growth factor 2; FGF-2; Basic fibroblast |
| | growth factor; bFGF; Heparin-binding growth factor 2; |
| | HBGF-2 |
| Mol Mass | 16.3 KDa |
| AP Mol Mass | 17 KDa, reducing conditions |
| Purity | Greater than 95% as determined by reducing |
| | SDS-PAGE. |
| Endotoxin | Less than 0.001 ng/ μ g (0.01 EU/ μ g) as determined by |
| | LAL test. |
| FORMULATION | Lyophilized from a 0.2 μ m filtered solution of 20mM |
| | Tris-HCl, 100mM NaCl, 0.02% Tween80, pH7.5. |



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com 23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.Q



RECONSTITUTION Always centrifuge tubes before opening.Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. SHIPPING The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. Lyophilized protein should be stored at < -20 ° C, STORAGE though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. BACKGROUND FGF-basic is a members of the Fibroblast Growth Factors (FGFs) family. The family constitutes a large family of proteins involved in many aspects of development including cell proliferation, growth, and differentiation. They act on several cell types to



+86-27-59760950

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.Q


ELKbio@ELKbiotech.com

regulate diverse physiologic functions including

www.elkbiotech.com



angiogenesis, cell growth, pattern formation, embryonic development, metabolic regulation, cell migration, neurotrophic effects, and tissue repair. FGF-basic is a non-glycosylated heparin binding growth factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland liver, monocytes, epithelial cells and endothelial cells.



SDS-PAGE



+86-27-59760950 ELKbio@ELKbiotech.com

n.com www.elk

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C