



Recombinant Mouse TfR (N-8His)

Catalog #	EPT081
Expression Host	Human Cells
DESCRIPTION	Recombinant Mouse Transferrin Receptor Protein 1 is produced by our Mammalian expression system and the target gene encoding Cys89-Phe763 is expressed with a 8His tag at the N-terminus.
Accession	Q62351
Synonyms	Transferrin receptor protein 1; TR; TfR; TfR1; Trfr; CD71; Tfrc
Mol Mass	77 KDa
AP Mol Mass	90 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Endotoxin	Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
FORMULATION	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, 5% Trehalose, 5% Mannitol, 0.01% tween80, pH 7.4.





RECONSTITUTION

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100 μ g/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at $< -20^{\circ}\text{C}$, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at $4-7^{\circ}\text{C}$ for 2-7 days.

Aliquots of reconstituted samples are stable at $< -20^{\circ}\text{C}$ for 3 months.

BACKGROUND


Transferrin receptor protein 1 (TFRC) belongs to the peptidase M28 family that is synthesized as a 172 amino acid (aa). TFRC regulated by cellular iron levels through binding of the iron regulatory proteins, IRP1 and IRP2, to iron-responsive elements in the 3'-UTR. It binds one transferrin or HFE molecule per subunit and





binds the HLA class II histocompatibility antigen, DR1. It Interacts with SH3BP3 and STEAP3, facilitates TFRC endocytosis in erythroid precursor cells. Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system. A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. It positively regulates T and B cell proliferation through iron uptake.

SDS-PAGE





ELK Biotechnology



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C