

Recombinant Human IFN gamma

Catalog # EPT166

Expression Host Human Cells

DESCRIPTION Recombinant Human Interferon Gamma is produced

by our Mammalian expression system and the target

gene encoding Gln24-Gln166 is expressed.

Accession P01579

Synonyms Interferon Gamma; IFN-Gamma; Immune Interferon;

IFNG

Mol Mass 16.8 KDa

AP Mol Mass 16&20&25 KDa, reducing conditions

Purity Greater than 95% as determined by reducing

SDS-PAGE.

Endotoxin Less than 0.1 ng/μg (1 EU/μg) as determined by LAL

test.

FORMULATION Lyophilized from a 0.2 µm filtered solution of 20mM

PB, 4% Mannitol, 2% Sucrose, 0.02% Tween80, pH7.4.

RECONSTITUTION Always centrifuge tubes before opening.Do not mix by

vortex or pipetting.



www.elkbiotech.com



It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at < -20 ° C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days.

Aliquots of reconstituted samples are stable at < -20° C for 3 months.


BACKGROUND

IFN γ is the major interferon produced by mitogenically or antigenically stimulated lymphocytes. It is structurally different from type I interferon and its major activity is immunoregulation. It has been implicated in the expression of class II histocompatibility antigens in cells that do not normally produce them, leading to autoimmune disease. Interferon gamma is produced mainly





byT-cells and natural killer cells activated by antigens, mitogens, or alloantigens. It is produced by lymphocytes expressing the surface antigens CD4 and CD8. IFNy synthesis is induced by IL-2, FGF-basic, and EGF.



SDS-PAGE



+86-27-59760950