



Recombinant Human LGALS3 (C-6His)

Catalog #	EPT190
Expression Host	Human Cells
DESCRIPTION	Recombinant Human Galectin-3 is produced by our Mammalian expression system and the target gene encoding Ala2-Ile250 is expressed with a 6His tag at the C-terminus.
Accession	AAH53667.1
Synonyms	Galectin-3; Gal-3; 35 kDa Lectin; Carbohydrate-Binding Protein 35; CBP 35; Galactose-Specific Lectin 3; Galactoside-Binding Protein; GALBP; IgE-Binding Protein; L-31; Laminin-Binding Protein; Lectin L-29; Mac-2 Antigen; LGALS3; MAC2
Mol Mass	27.2 KDa
AP Mol Mass	35 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Endotoxin	Less than 0.1 ng/μg (1 EU/μg) as determined by LAL





test.

FORMULATION

Lyophilized from a 0.2 μ m filtered solution of PBS, 3mM DTT, pH 7.4.

RECONSTITUTION

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100 μ g/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at $< -20^{\circ}$ C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7 $^{\circ}$ C for 2-7 days.

Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.

BACKGROUND

Galectin-3(LGALS3) is also known as Galactose-specific lectin 3, Mac-2 antigen, Carbohydrate-binding protein 35, Laminin-binding





protein and Galactoside-binding protein. LGALS3 is highly expressed in early stages of papillary carcinoma, and lowly during tumor progression. LGALS3 is probably forms homo- or heterodimers and secreted by a non-classical secretory pathway and associates with the cell surface. LGALS3 plays an important role during the acquisition of vasculogenic mimicry and angiogenic properties. LGLAS3 takes part in an immune regulator to inhibit T-cell immune responses and promote tumor growth, as a result providing a new mechanism for tumor immune tolerance.

SDS-PAGE

