

Recombinant Human TGF-beta 2

Catalog # EPT195

Expression Host Human Cells

DESCRIPTION Recombinant Human Transforming Growth Factor

Beta 2 is produced by our Mammalian expression

system and the target gene encoding Ala303-Ser414 is

expressed.

Accession P61812

Synonyms Transforming growth factor beta-2; TGFB2; Polyergin;

G-TSF; Glioblastoma-derived T-cell suppressor factor;

Cetermin; BSC-1 cell growth inhibitor; TGF-beta-2

Mol Mass 12.7 KDa

AP Mol Mass 12 KDa, reducing conditions

Purity Greater than 95% as determined by reducing

SDS-PAGE.

Endotoxin Less than 0.001 ng/μg (0.01 EU/μg) as determined by

LAL test.

FORMULATION Lyophilized from a 0.2 µm filtered solution of 4mM

HCI.



www.elkbiotech.com



RECONSTITUTION

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at < -20 $^{\circ}$ C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days.

Aliquots of reconstituted samples are stable at < -20° C for 3 months.

BACKGROUND

Transforming growth factor beta-2 (TGF- β 2) is a secreted protein which belongs to the TGF-beta family. It is known as a cytokine that performs many cellular functions and has a vital role during embryonic development. The precursor is cleaved into mature TGF-beta-2 and LAP, which remains non-covalently





linked to mature TGF-beta-2 rendering it inactive. It is an extracellular glycosylated protein. It is known to suppress the effects of interleukin dependent T-cell tumors. Defects in TGFB2 may be a cause of non-syndromic aortic disease (NSAD).

