



# Recombinant Human NAMPT (N-6His)

<b>Catalog #</b>	EPT199
<b>Expression Host</b>	E.coli
<b>DESCRIPTION</b>	Recombinant Human Pre-B-Cell Colony-Enhancing Factor 1 is produced by our E.coli expression system and the target gene encoding Met1-His491 is expressed with a 6His tag at the N-terminus.
<b>Accession</b>	P43490
<b>Synonyms</b>	Pre-B cell-enhancing factor; Nicotinamide phosphoribosyltransferase; NAmPRTase; Nampt; Pre-B-cell colony-enhancing factor 1; Visfatin; NAMPT; PBEF; PBEF1
<b>Mol Mass</b>	57 KDa
<b>AP Mol Mass</b>	55 KDa, reducing conditions
<b>Purity</b>	Greater than 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
<b>FORMULATION</b>	Lyophilized from a 0.2 μm filtered solution of 20mM





HEPES, 150mM NaCl, pH 8.0.

## RECONSTITUTION

Always centrifuge tubes before opening. Do not mix by vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

## STORAGE

Lyophilized protein should be stored at  $< -20^{\circ}\text{C}$ , though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at  $4-7^{\circ}\text{C}$  for 2-7 days.

Aliquots of reconstituted samples are stable at  $< -20^{\circ}\text{C}$  for 3 months.

## BACKGROUND

Pre-B cell colony enhancing factor (PBEF) was originally identified as a cytokine that potentiated the clonal expansion and differentiation of pre-B cells, but it is also acknowledged to be the ubiquitous intracellular enzyme nicotinamide





phosphoribosyltransferase (NAMPT) and the adipokine “visfatin” . PBEF is constitutively expressed in the fetal membranes where its greatest expression is in the amnion. It has intracellular and extracellular forms. Most of the intracellular functions of PBEF are due to its role as a Nampt which can induce angiogenesis through upregulation of VEGF and VEGFR and secretion of MCP-1. Extracellular PBEF has been shown to increase inflammatory cytokines, such as TNF- $\alpha$ , IL-1  $\beta$ , IL-16, and TGF-  $\beta$  1. PBEF also increases the production of IL-6, TNF-  $\alpha$  , and IL-1  $\beta$  in CD14+ monocytes, macrophages, and dendritic cells, enhances the effectiveness of T cells.





**ELK Biotechnology**

kDa MK R


120

90

60

40

30



**SDS-PAGE**



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C