

Recombinant Human TPBG (C-6His)

Catalog # EPT261

Expression Host Human Cells

DESCRIPTION Recombinant Human Trophoblast Glycoprotein is

produced by our Mammalian expression system and

the target gene encoding Ser32-Ser355 is expressed

with a 6His tag at the C-terminus.

Accession Q13641

Synonyms 5T4 Protein, Human;5T4AG Protein, Human;M6P1

Protein, Human; WAIF1 Protein, Human

Mol Mass 36.1 KDa

AP Mol Mass 55-80 KDa, reducing conditions

Purity Greater than 95% as determined by reducing

SDS-PAGE.

Endotoxin Less than 0.1 ng/μg (1 EU/μg) as determined by LAL

test.

FORMULATION Lyophilized from a 0.2 µm filtered solution of PBS,

pH7.4.

RECONSTITUTION Always centrifuge tubes before opening.Do not mix by



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



vortex or pipetting.

It is not recommended to reconstitute to a concentration less than 100µg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at < -20 ° C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days.

Aliquots of reconstituted samples are stable at < -20° C for 3 months.

BACKGROUND

Trophoblast glycoprotein, also known as TPBG and 5T4, it belongs to the LRR family of proteins. The C-terminus of LRR N-terminal cap (LRRNT) and LRR 1 are essential for the inhibition of the Wnt signaling pathway. TPBG may function as an inhibitor of Wnt/beta-catenin signaling by indirectly interacting with LRP6 and blocking Wnt3a-dependent LRP6





internalization. It expressed by all types of trophoblasts as early as 9 weeks of development. In adult tissues, the expression is limited to a few epithelial cell types but is found on a variety of carcinoma.



+86-27-59760950